

# Seiberg Duality in Matrix Model

---

**Bo Feng**

*Institute for Advanced Study  
Einstein Drive,  
Princeton, New Jersey, 08540*

**ABSTRACT:** In this paper, we use the matrix model to show the Seiberg duality in the case of complete mass deformation.

**KEYWORDS:** Seiberg Duality, Matrix Model.

**Motivation** Since Dijkgraaf and Vafa gave the matrix model conjecture [1, 2, 3], there are a lot of related works to check, prove and generalize this conjecture, see [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25]. Especially in [15] and [23], a field theory argument has been given to show why the calculation of exact lower energy superpotential can be reduced to integration in the corresponding matrix model.

The matrix model conjecture holds for more general cases although the primary focus is still the theory with adjoint fields. The generalization to fundamental fields has been discussed in [14, 13, 16, 18, 19]. These kinds of generalization are desired because most applications in field theory will have fundamental matters. One particular interesting application is to see the Seiberg duality [26] in the matrix model.

The standard example of Seiberg dual pair is following. At one side, we have electric theory  $SU(N_c)$  with  $N_f$  flavors  $Q_j, \tilde{Q}_j$  and no superpotential. At another side, we have magnetic theory  $SU(N_f - N_c)$  with  $N_f$  flavors  $q_j, \tilde{q}_j$ , meson field  $X_i^j$  and superpotential

$$W_{mag} = \frac{1}{\mu} X_i^j q_j \tilde{q}^i. \quad (1)$$

where  $\mu$  is a dynamical scale [27] (equation (3.125)). In general case, these two theories will have flat directions in the moduli space, so when we try to do the matrix model integration, we must take care of the zero modes as did by Berenstein[11]. To avoid the complexity, we can deform above theories by adding mass terms. For example, in the electric theory we add superpotential with non-degenerated mass matrix<sup>1</sup>

$$W_{elec} = \sum_{j=1}^{N_f} Q_j m_j \tilde{Q}^j \quad (2)$$

Since we give all flavors nonzero mass in the electric theory, the IR field theory will be pure super Yang-Mills theory and we know the exact effective action as

$$W_{YM} = N_c (\hat{\Lambda}^{3N_c})^{\frac{1}{N_c}} \quad (3)$$

where  $\hat{\Lambda}$  is the dynamical scale at IR and related to dynamical scale  $\Lambda$  at UV by

$$\hat{\Lambda}^{3N_c} = \det(m) \Lambda^{3N_c - N_f} \quad (4)$$

The deformation in the electric theory will induce the corresponding deformation in magnetic theory as

$$W_{mag} = \frac{1}{\mu} X_i^j q_j \tilde{q}^i + \text{tr}(Xm) \quad (5)$$

---

<sup>1</sup>We can always redefine the field to bring the mass matrix into diagonal form.

**Matrix integration** Now we will do the matrix integration for both electric and magnetic theories. Let us do the electric theory first. The matrix model is

$$\frac{1}{\text{Vol}(U(N))} \int \prod_{j=1}^{N_f} dQ_j dQ_j^\dagger e^{-\frac{1}{g_s} \sum_{j=1}^{N_f} Q_j m_j Q_j^\dagger} \quad (6)$$

where every  $Q$  is a  $M$  component vector and  $\tilde{Q}$  in the field theory has been treated as a conjugate of  $Q$  [2]. The part of flavor integration is

$$\begin{aligned} \int \prod_{j=1}^{N_f} dQ_j dQ_j^\dagger e^{-\frac{1}{g_s} \sum_{j=1}^{N_f} Q_j m_j Q_j^\dagger} &= \prod_{j=1}^{N_f} \left( \frac{\pi g_s}{m_j} \right)^M \\ &= (\pi g_s)^{MN_f} (\det(m))^{-M} = e^{MN_f \log(\pi g_s) - M \log(\det(m))} \end{aligned}$$

Except that, there is also the volume factor  $e^{\frac{M^2}{2} \log M}$ . Adding everything together we get

$$\begin{aligned} &\exp\left(\frac{M^2}{2} \log M + MN_f \log(\pi g_s) - M \log(\det(m))\right) \\ &= \exp\left(\frac{1}{g_s^2} \frac{S^2}{2} \log \frac{S}{g_s} + \frac{1}{g_s} [SN_f \log(\pi g_s) - S \log(\det(m))]\right) \\ &\equiv \exp\left(\frac{1}{g_s^2} \mathcal{F}_{\chi=2} + \frac{1}{g_s} \mathcal{F}_{\chi=1}\right) \end{aligned}$$

where we have grouped all term according to the genus expansion. From the corresponds given in [3, 13], we should identify

$$-W_{elec} = N_c \frac{\partial(\frac{S^2}{2} \log \frac{S}{g_s})}{\partial S} + [SN_f \log(\pi g_s) - S \log(\det(m))] \quad (7)$$

In the matrix model,  $S, M$  are dimensionless number, but in field theory,  $S$  is dimension 3. To match the field theory result, we need to replace  $g_s$  by some proper dimensional number heuristically. For example, we need to replace  $\frac{S}{g_s}$  to  $\frac{S}{\Lambda^3 e^{3/2}}$  to reproduce the well known Veneziano-Yankielowicz term. Same heuristic argument tell us that  $\pi g_s$  in the second term must be replaced by dimension one number which can be chosen naturally as  $\Lambda$ . Doing these replacement we get

$$\begin{aligned} -W_{elec} &= N_c \frac{\partial(\frac{S^2}{2} \log \frac{S}{\Lambda^3 e^{3/2}})}{\partial S} + [SN_f \log(\Lambda) - S \log(\det(m))] \\ &= N_c [S \log \frac{S}{\Lambda^3} - S] + [SN_f \log(\Lambda) - S \log(\det(m))] \\ &= N_c [S \log(\frac{S}{(\Lambda^{3N_c - N_f} \det(m))^{\frac{1}{N_c}}}) - S] \end{aligned}$$

Minimized it we get  $S = (\Lambda^{3N_c - N_f} \det(m))^{\frac{1}{N_c}}$  and

$$W_{elec} = N_c (\hat{\Lambda}^{3N_c})^{\frac{1}{N_c}} = N_c (\Lambda^{3N_c - N_f} \det(m))^{\frac{1}{N_c}} \quad (8)$$

which is the famous Affleck-Dine-Seiberg superpotential for pure  $U(N)$  Yang-Mills gauge theory. Notice that the equation (4) is naturally shown in the matrix model. Furthermore, in our calculation, we keep  $N_f$  fixed while taking  $M \rightarrow \infty$ .

Now we do the matrix model integration for the magnetic theory

$$\frac{1}{\text{Vol}(U(N))} \int dX \prod_j dq_j dq_j^\dagger \exp\left(\frac{-1}{g_s} [\text{tr}(mX) + \sum_{i,j=1}^{N_f} \frac{1}{\mu} X_i^j q_j q^{\dagger i}]\right) \quad (9)$$

where  $X$  is  $N_f \times N_f$  matrix and  $q$  is  $N$  component vector. Doing the  $X$  integration first, we get a delta-function

$$\delta\left(m + \frac{q_j q^{\dagger i}}{\mu}\right) = \mu^{N_f^2} \delta(\mu m + q_j q^{\dagger i}) \quad (10)$$

Now the remainder part is exact the integration given in [19] (equation (6) with  $-\mu m$  taking the place of  $X$  there. Furthermore, since we always take large  $N$  limit with fixed  $N_f$ , there is not problem to apply their result.), so we just cite their result

$$W_{mag} = (\widetilde{N}_c - N_f) \left( \frac{\widetilde{\Lambda}^{3\widetilde{N}_c - N_f}}{\det(-\mu m)} \right)^{\frac{1}{\widetilde{N}_c - N_f}} \quad (11)$$

where we use tilde to emphasize that it is in the magnetic theory. Using the result  $\widetilde{N}_c - N_f = -N_c$ ,  $\det(-\mu m) = (-)^{N_f} \mu^{N_f} \det(m)$  and

$$\Lambda^{3N_c - N_f} \widetilde{\Lambda}^{3\widetilde{N}_c - N_f} = (-)^{N_f - N_c} \mu^{N_f}, \quad (12)$$

which can be found, for example, in [27]. we get immediately

$$W_{mag} = N_c (\Lambda^{3N_c - N_f} \det(m))^{\frac{1}{N_c}} \quad (13)$$

where the  $-$  sign in front of (11) is canceled exactly by factor  $(-)^{N_c}$ . Comparing (8) and (13) we see that they are same. Thus we give an example to show how the matrix model can see the Seiberg duality.

We must emphasize that we did not really derive the Seiberg duality from the matrix model because the Seiberg duality holds in general situation even without any mass deformation. We feel that to address the full Seiberg duality, we must understand how to do the matrix integration when there is flat directions in moduli space along the line [11]. Obvious, work should be generalized other generalized Seiberg duality, for example, the toric duality addressed in [28, 29, 30, 31, 32, 33, 34].

**Acknowledgements** We deeply thank Nathan Seiberg for suggesting the mass deformation. We also likes to thank Vijay Balasubramanian, David Berenstein, Freddy Cachazo, Joshua Erlich, Yang-Hui He, Mxing Huang, Vishnu Jejjala and Asad Naqvi for fruitful discussion. This research is supported under the IAS grant **PHY-0070928**.

## References

- [1] R. Dijkgraaf and C. Vafa, “Matrix models, topological strings, and supersymmetric gauge theories,” Nucl. Phys. B **644**, 3 (2002) [arXiv:hep-th/0206255].
- [2] R. Dijkgraaf and C. Vafa, “On geometry and matrix models,” Nucl. Phys. B **644**, 21 (2002) [arXiv:hep-th/0207106].
- [3] R. Dijkgraaf and C. Vafa, “A perturbative window into non-perturbative physics,” arXiv:hep-th/0208048.
- [4] N. Dorey, T. J. Hollowood, S. P. Kumar and A. Sinkovics, “Massive vacua of  $N = 1^*$  theory and S-duality from matrix models,” arXiv:hep-th/0209099.
- [5] N. Dorey, T. J. Hollowood, S. Prem Kumar and A. Sinkovics, “Exact superpotentials from matrix models,” arXiv:hep-th/0209089.
- [6] N. Dorey, T. J. Hollowood and S. P. Kumar, “S-duality of the Leigh-Strassler deformation via matrix models,” arXiv:hep-th/0210239.
- [7] L. Chekhov and A. Mironov, “Matrix models vs. Seiberg-Witten/Whitham theories,” arXiv:hep-th/0209085.
- [8] F. Ferrari, “On exact superpotentials in confining vacua,” arXiv:hep-th/0210135.
- [9] H. Fuji and Y. Ookouchi, “Comments on effective superpotentials via matrix models,” arXiv:hep-th/0210148.
- [10] R. Dijkgraaf, S. Gukov, V. A. Kazakov and C. Vafa, “Perturbative analysis of gauged matrix models,” arXiv:hep-th/0210238.
- [11] D. Berenstein, “Quantum moduli spaces from matrix models,” arXiv:hep-th/0210183.
- [12] A. Gorsky, “Konishi anomaly and  $N = 1$  effective superpotentials from matrix models,” arXiv:hep-th/0210281.
- [13] R. Argurio, V. L. Campos, G. Ferretti and R. Heise, “Exact superpotentials for theories with flavors via a matrix integral,” arXiv:hep-th/0210291.
- [14] J. McGreevy, “Adding flavor to Dijkgraaf-Vafa,” arXiv:hep-th/0211009.
- [15] R. Dijkgraaf, M. T. Grisaru, C. S. Lam, C. Vafa and D. Zanon, “Perturbative Computation of Glueball Superpotentials,” arXiv:hep-th/0211017.
- [16] H. Suzuki, “Perturbative Derivation of Exact Superpotential for Meson Fields from Matrix Theories with One Flavour,” arXiv:hep-th/0211052.
- [17] F. Ferrari, “Quantum parameter space and double scaling limits in  $N=1$  super Yang-Mills theory,” arXiv:hep-th/0211069.
- [18] I. Bena and R. Roiban, “Exact superpotentials in  $N=1$  theories with flavor and their matrix model formulation,” arXiv:hep-th/0211075.

- [19] Y. Demasure and R. A. Janik, “Effective matter superpotentials from Wishart random matrices,” arXiv:hep-th/0211082.
- [20] M. Aganagic, A. Klemm, M. Marino and C. Vafa, “Matrix Model as a Mirror of Chern-Simons Theory,” arXiv:hep-th/0211098.
- [21] R. Gopakumar, “ $\mathcal{N} = 1$  Theories and a Geometric Master Field,” arXiv:hep-th/0211100.
- [22] S. Naculich, H. Schnitzer and N. Wyllard, “The  $N = 2$   $U(N)$  gauge theory prepotential and periods from a perturbative matrix model calculation,” arXiv:hep-th/0211123.
- [23] F. Cachazo, M. R. Douglas, N. Seiberg and E. Witten, “Chiral Rings and Anomalies in Supersymmetric Gauge Theory,” arXiv:hep-th/0211170.
- [24] R. Dijkgraaf, A. Neitzke and C. Vafa, “Large  $N$  Strong Coupling Dynamics in Non-Supersymmetric Orbifold Field Theories,” arXiv:hep-th/0211194.
- [25] Y. Tachikawa, “Derivation of the Konishi anomaly relation from Dijkgraaf-Vafa with (Bi-)fundamental matters,” arXiv:hep-th/0211189.
- [26] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” Nucl. Phys. B **435**, 129 (1995) [arXiv:hep-th/9411149].
- [27] P. C. Argyres, “Introduction to supersymmetry”, Course note.
- [28] B. Feng, A. Hanany and Y. H. He, “D-brane gauge theories from toric singularities and toric duality,” Nucl. Phys. B **595**, 165 (2001) [arXiv:hep-th/0003085].
- [29] B. Feng, A. Hanany and Y. H. He, “Phase structure of D-brane gauge theories and toric duality,” JHEP **0108**, 040 (2001) [arXiv:hep-th/0104259].
- [30] B. Feng, A. Hanany, Y. H. He and A. M. Uranga, “Toric duality as Seiberg duality and brane diamonds,” JHEP **0112**, 035 (2001) [arXiv:hep-th/0109063].
- [31] B. Feng, S. Franco, A. Hanany and Y. H. He, “Symmetries of toric duality,” arXiv:hep-th/0205144.
- [32] F. Cachazo, B. Fiol, K. A. Intriligator, S. Katz and C. Vafa, “A geometric unification of dualities,” Nucl. Phys. B **628**, 3 (2002) [arXiv:hep-th/0110028].
- [33] C. E. Beasley and M. R. Plesser, “Toric duality is Seiberg duality,” JHEP **0112**, 001 (2001) [arXiv:hep-th/0109053].
- [34] D. Berenstein and M. R. Douglas, “Seiberg duality for quiver gauge theories,” arXiv:hep-th/0207027.